

# IN-BEAM GAMMA-RAY SPECTROSCOPY OF TARGET FRAGMENTATION<sup>\*</sup>

E. Rodriguez-Vieitez<sup>a,b)</sup>, R. M. Clark<sup>a)</sup>, M. Cromaz<sup>a)</sup>, M. A. Deleplanque<sup>a)</sup>, M. Descovich<sup>a)</sup>,  
R. M. Diamond<sup>a)</sup>, P. Fallon<sup>a)</sup>, A. Goergen<sup>a,c)</sup>, G. J. Lane<sup>a,d)</sup>, I. Y. Lee<sup>a)</sup>, A. O. Macchiavelli<sup>a)</sup>,  
H. Mahmud<sup>a)</sup>, S. G. Prussin<sup>b)</sup>, F. S. Stephens<sup>a)</sup>, C. E. Svensson<sup>a,e)</sup>, K. Vetter<sup>a,f)</sup>, D. Ward<sup>a)</sup>

<sup>a)</sup>Lawrence Berkeley National Laboratory, Berkeley, CA 94720, USA

<sup>b)</sup>University of California, Berkeley, CA 94720, USA

<sup>c)</sup>CEA Saclay, F-91191 Gif-sur-Yvette, France

<sup>d)</sup>Australian National University, Canberra, Australia

<sup>e)</sup>University of Guelph, Guelph, Ontario N1G2W1, Canada

<sup>f)</sup>Lawrence Livermore National Laboratory, Livermore, CA 94550, USA

Fragmentation reactions at energies  $\sim E/A > 50$  MeV produce neutron-rich nuclei but leave little angular momentum in the residues. In this work we have examined the product distribution and angular momentum input for a  $^{12}\text{C}$  beam at 30 MeV/A on a thick (40 mg/cm<sup>2</sup>) target of  $^{51}\text{V}$ , testing the feasibility of in-beam gamma-ray spectroscopy. With this technique it might be possible to study neutron-rich nuclei to moderate spins, complementing the now traditional beam fragmentation studies. The main advantage of the target fragmentation setup is that, for lifetimes longer than the stopping time ( $\sim 1$  ps), gamma rays are not Doppler broadened. The experiment was performed with the Gammasphere array and the beam delivered by the 88-inch cyclotron at LBNL.

We will present the results of this study in which some 70 different isotopes from F to Fe ( $Z=9-26$ ) have been identified and new excited states found. The experimental isotopic yields, such as those shown in Figure 1, cannot be reproduced by LISE calculations [1], which are based on the abrasion-ablation model of fragmentation. Also, the maximum angular momentum of 12 that is observed experimentally is greater than that expected from a model [2] based on fragmentation. At the low energy of this experiment (30 MeV/A) a wide variety of reactions may be taking place including fragmentation, transfer, deep-inelastic, incomplete fusion, etc. More comprehensive reaction models, such as those using antisymmetrized molecular dynamics [3], could be appropriate and may be investigated in the future.

<sup>\*</sup>This work has been supported by the U.S. D.o.E. under Contract No. DE-AC03-76SF00098

- [1] D. Bazin *et al.*, Nuclear Instruments and Methods in Physics Research A **482**, 307-327 (2002).
- [2] M. P. Pfützner *et al.*, Physical Review C **65**(6), 064604 (2002).
- [3] A. Ono *et al.*, Physical Review Letters **68**(19), 2898 (1992).




Figure 1. Selected yields observed in the experiment (30 MeV/A  $^{12}\text{C}$  beam on a 40 mg/cm<sup>2</sup>  $^{51}\text{V}$  target).